


Mortality in Bay County

2012 Bay County Community Health Assessment Data




Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010

The majority of deaths (**349**) among adults in Bay County were due to **Heart Disease**.


Cancer followed closely causing **269** deaths.

Of the 349 deaths due to heart disease, **255** of them were among people **75 years of age or older**.

More women died from heart disease than men did.



Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010




Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010

Cancer was responsible for more deaths among **45-74** year olds.

The number of deaths caused by cancer among **males** and **females** are relatively equal.

Mortality in Bay County Continued


2012 Bay County Community Health Assessment Data



Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010

Chronic Lower Respiratory Disease accounted for a total of **60 deaths** among Bay County adults.


Of those, the **majority of deaths** occurred among **females 75 years and older**.



Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010

A total of **53 deaths** from **strokes** occurred among Bay County adults in 2009.

Of those, the **majority of deaths** occurred among **females 75 years and older**.



Source: Michigan Behavioral Risk Factor Surveillance Survey. 2010

Heart disease and cancer are the two leading causes of death for Bay County Residents. These deaths attribute to a **total of 3,929 years of potential life lost** for persons under 75 years of age.